# Exhibit 307 (Filed Under Seal)

To: Creech, Kristen[Kristen.Creech@actavis.com]

From: Actavis Mase 1:15-cv-07488-CM-RWL Document 467-92 Filed 12/22/17 Page 2 of 6

Sent: Fri 5/22/2015 8:07:40 PM Importance: Normal

Subject: Actavis Confirms Appeals Court Ruling Requiring Continued Distribution of NAMENDA IR

Received: Fri 5/22/2015 8:07:43 PM

Actavis plc has added a press release to its Investor Relations Web site.

Title: Actavis Confirms Appeals Court Ruling Requiring Continued Distribution of NAMENDA IR

Date(s): May. 22, 2015 4:05 PM

For a complete listing of our news releases, please click here

DUBLIN, May 22, 2015 /PRNewswire/ — Actavis pic (NYSE: ACT) today confirmed that a panel of the U.S. Court of Appeals for the Second Circuit has issued a ruling upholding a December 15, 2014 preliminary injunction requiring the Company to continue distribution of NAMENDA® (memantine HCI) immediate-release tablets.

Logo - http://photos.prnewswire.com/pmh/20130124/NY47381LOGO

"While we are disappointed by the Court's decision to uphold this ruling, we intend to continue our strong efforts to convey the significant benefits of NAMENDA XR® to physicians, patients and caregivers," said Brent Saunders, CEO and President of Actavis. "Patient demand for NAMENDA XR® is currently trending at more than 50 percent of the total product line's days of therapy and growing, underscoring the strong physician, patient and caregiver demand for our once-daily product."

We have also recently launched once-daily NAMZARIC®, a fixed-dose combination of NAMENDA XR® and donepezil that provides another treatment option for patients with moderate to severe Alzheimer's disease. Since the launch of NAMENDA XR® in 2013, the two medications, NAMENDA XR® and donepezil, have been commonly prescribed in combination with one another to treat the symptoms of moderate to severe Alzheimer's disease. NAMZARIC® offers an option with the benefits of both treatments, while reducing the number of pills a patient and their caregivers need to administer each day, to treat this disease."

Actavis noted that the Company will continue to manage sales and R&D expenses to ensure that the Appeals court's decision will have minimal to no impact on its 2015 NAMENDA® franchise contribution to earnings and longer term company earnings aspirations.

About NAMENDA XR®

NAMENDA XR® (memantine HCI) extended release capsules are a higher dose, once-daily formulation of NAMENDA® immediate release indicated for the treatment of moderate to severe dementia of the Alzheimer's type. Its mechanism of action focuses on the glutamate pathway, a target for the treatment of Alzheimer's disease. The efficacy and safety of NAMENDA XR® was established in a 24 week, randomized, double-blind, placebo-controlled trial of 677 outpatients on a stable dose of acetylcholinesterase inhibitors (AChEI).

NAMENDA XR® 28 mg plus an AChEl demonstrated statistically significant improvement in cognition and global function compared to placebo plus an AChEl, Cognition was measured by the Severe Impairment Battery Scale (2.6 unit mean difference). Global

function was measured by the Clinician's Interest 1.15-cv-07488-C	erview-Based Impressi M-RWL Documer	on of Change Scale ( at 467-92 Filed 1	0.3. unit mean difference 2/22/17 Page 3 of	6
Those is an evidence that MANENIDA VIDE a	ne on AChEl neovocia a	e elomo tha madachida		alata ilitela
There is no evidence that NAMENDA XR® c Alzheimer's disease.	an Achei prevents o	r slows the underlying	oisease process at pati	ents with
		n de la proposición de la composición de la proposición de la proposición de la proposición de la proposición La proposición de la		u. 12 - Tagang angganggang
	1 [독급] [ [ [ [ [ ]			
Dosing and Administration	영화 기원을 감독되렴		) (15== - 1) - (24)	
And the second s				ik Verioù die Serie i
				스케이 이 사람이 함께 그는 그는 사람들이 함께
The recommended starting dose of increased in 7 mg increases is one week and only if the increase.	increments to 28 mg operations dose has been	once daily. The minim n well tolerated, The r	um recommended interv naximum recommended	al between dose dose is 28 mg onc
It is recommended that a patient who XR® 28 mg once-daily capsules the dathe comparative efficacy of these 2 recommended that a patient will tablets be switched to NAMENDA XR	lay following the last do gimens. h severe renal impairm	se of a 10 mg NAMEI ent who is on a regim	NDA® tablet. There is no en of 5 mg twice daily of	study addressing
Special Populations			기를 위치점한 기가 하는 기자 기 사람들은 기계를 당된 기가 보고 하다	
[속이 기계속이라고 한국의 공항] (2017)				
			= =====================================	
NAMENDA XR® should be administ     A target dose of 14 mg/day is recombased on the Cockcroft-Gault equation	nmended in patients wit			ce of 5-29 mL/min,
		***	=	•
IMPORTANT SAFETY INFORMATION				
Contraindications				
	,			
NAMENDA XR is contraindicated in	patients with known hy	persensitivity to mem	antine hydrochloride or t	o any excipients
used in the formulation.	ruklerkkuk d			
				과 그는 바둑
Warnings and Precautions				
				SHE STATE
NAMENDA XR should be used with clinical state of the patient). Alkaline unincreased plasma levels and a possible NAMENDA XR has not been system.	rine conditions may de le increase in adverse (	crease the urinary elireffects.	nination of memantine, n	diet, drugs and the esulting in
이 등이 있다는 그는 이 사람들은 독자를 하고				
			rough on, her a section of	1995 - Ar. 54
Adverse Reactions			n nga <u>s</u> a S <u>elatata</u> sa Selata . Selata sa Selata	
	on de la ferio de <del>la c</del> elo filo La companya de la co			
• The most commonly observed advan	rea reactions cono in a	ationts administered b	NAMENDA VD 129 mold	au) in a controllad
The most commonly observed adve- clinical trial, defined as those occurring than placebo were headache (6% vs.5	g at a frequency of at le	east 5% in the NAME	NDA XR group and at a l	nigher frequency
grand to the property of the second of the s				•

Case	1:15-cv-07488-CM-RWL	Document 467-92	Filed 12/22/17	Page 4 of 6	
Drug Interactions					_ :-
No data-data ir	nteraction studies have been co	onducted with NAMENDA	XR specifically The	e combined use of l	NAMENDA
XR with other NN	MDA antagonists (amantadine, I be approached with caution				
	ရုံရေး သည် ရှေးသည် မိုင်ငံ ရုံရေးကြွေ				
Please visit www.Name	endaXR.com for more informati	on and full prescribing in	formation.		
About NAMZARICTM				- , , , , , , , , , , , , , , , , , , ,	•
ABOUT HAMEAING		•			
	en de la companya de La companya de la co	re Signatur (1988)		Bayer Bayer Santago a la Sena de la Caracteria	
donepezii hydrochloride (memantine HCI extend severe renal impairmer XR®, which is indicated in ARICEPT® (donepe Actavis and Adamas co	laily, fixed-dose combination of e, an acetylcholinesterase inhib ded-release/donepezil HCl) and nt. Memantine hydrochloride ex d for the treatment of moderate zil hydrochloride), which is indic ollaborated on the development s, while Adamas will retain excl	oitor. NAMZARIC will be a last 14/10mg (memantine Hatended-release is the act to severe dementia of the cated for the treatment of the fixed-dose combined.	available in two dosaged extended-release/ ive ingredient in the control of the c	ge strengths, 28/10 donepezil HCl) for pour ently marketed lonepezil is the action of the Alzheimell have exclusive U.	mg patients with NAMENDA we ingredient er's type.
	s, witho Againas will retail exc	do to continuo cianzanon	ngnts outside of the		
					F 1
IMPORTANT SAFETY	INFORMATION				
			a filosofi University	ing tanggaran salah s Salah salah sa	
CONTRAINDICATION	<b>S</b>				
			Territoria. Territoria		
NAMES AND DESCRIPTIONS					مامام
	dicated in patients with known to be to any excipients used in the		nune nyarochionae, c	onepezii nydrociio	iikie,
					<u>-</u>
WARNINGS AND PRE	CAUTIONS				
		ng for a se <del>ta</del> a fu Magazar 2000 oo saa	palitera on A Resident	ouagren († 1920) Hongrafier	
Cardiovascular as bradycardia o cardiac conductic Peptic Ulcer Di symptoms of acti history of ulcer di Nausea and Vo been mild and tra Genitourinary Othe urinary elimin Seizures: Choliconvulsions. How	AMZARIC is likely to exaggerate Conditions: NAMZARIC may he heart block. Bradycardia or he on abnormalities. Is ease and Gastrointestinal Bleve or occult gastrointes	nave vagotonic effects on eart block may manifest in eding: Patients treated we eding, especially those a urrent nonsteroidal anti-in diarrhea, nausea, and vo- ng continued treatment. use bladder outflow obstances of plasma levels in increased plasma levels it hydrochloride, are belief be a manifestation of Al-	the sinoatrial and at a patients both with a ith NAMZARIC should be increased risk for different drugs (Nomiting, although in nuctions: Conditions of memantine, eved to have some patheimer's disease.	rioventricular nodes and without known used by monitored close eveloping ulcers, the ISAIDs).  The cases these effect that raise urine pH is otential to cause ge	inderlying sely for lose with a fects have may decreas eneralized
obstructive pulmo		a strong be breschised	mini care to patients	will a filology of ast	unita VI
ADVERSE REACTION	S		The second secon	s to the Sign of Later	<u> </u>

# Case 1:15-cv-07488-CM-RWL Document 467-92 Filed 12/22/17 Page 5 of 6

The most common adverse reactions, occurring at a frequency of at least 5% in patients taking memantine hydrochloride extended release 28 mg/day, and greater than placebo, were headache (6% vs 5%), diarrhea (5% vs 4%), and dizziness (5% vs 1%).

The most common adverse reactions, occurring at a frequency of at least 5% in patients taking donepezil, and at twice or more the rate of placebo, include diarrhea (10% vs 4%), anorexia (8% vs 4%), vomiting (8% vs 4%), nausea (6% vs 2%), and ecchymosis (5% vs 2%).

## DRUG INTERACTIONS

Alterations of urine pH toward the alkaline condition may lead to an accumulation of memantine with a possible increase in adverse reactions. NAMZARIC should be used with caution under conditions that may be associated with increased urine pH including alterations by diet and the clinical state of the patient.

The combined use of memantine hydrochloride with other NMDA antagonists (amantadine, ketamine, and dextromethorphan) has not been systematically evaluated and such use should be approached with caution.

Inhibitors of CYP450, 3A4 (e.g., ketoconazole) and 2D6 (e.g., quinidine), inhibit donepezil metabolism in vitro. Whether there is a clinical effect of quinidine is not known.

Inducers of CYP3A4 (e.g., phenytoin, carbamazepine, dexamethasone, rifampin, and phenobarbital) could increase the rate of elimination of done pezil.

Cholinesterase inhibitors, including donepezil hydrochloride, have the potential to interfere with the activity of anticholinergic medications.

### DOSAGE AND ADMINISTRATION

Patients stabilized on memantine hydrochloride (10 mg twice daily or 28 mg extended-release once daily and donepezil hydrochloride 10 mg) can be switched to NAMZARIC 28 mg/10 mg, taken once a day in the evening. Patients should start NAMZARIC the day following the last dose of memantine hydrochloride and donepezil hydrochloride administered separately. Patients with severe renal impairment (creatinine clearance 5-29 mL/min, based on the Cockcroft-Gault equation), stabilized on memantine hydrochloride (5 mg twice daily or 14 mg extended-release once daily) and donepezil hydrochloride 10 mg, can be switched to NAMZARIC 14 mg/10 mg, taken once daily.

### **About Actavis**

Actavis plc (NYSE: ACT), headquartered in Dublin, Ireland, is a unique, global pharmaceutical company and a leader in a new industry model - Growth Pharma. Actavis is focused on developing, manufacturing and commercializing innovative branded pharmaceuticals, high-quality generic and over-the-counter medicines and biologic products for patients around the world.

Actavis markets a portfolio of best-in-class products that provide valuable treatments for the central nervous system, eye care, medical aesthetics, gastroenterology, women's health, urology, cardiovascular and anti-infective therapeutic categories, and operates the world's third-largest global generics business, providing patients around the globe with increased access to affordable, high-quality medicines. Actavis is an industry leader in research and development, with one of the broadest development pipelines in the pharmaceutical industry and a leading position in the submission of generic product applications globally.

With commercial operations in approximately 100 countries, Actavis is committed to working with physicians, healthcare providers and patients to deliver innovative and meaningful treatments that help people around the world live longer, healthier lives.

Actavis intends to adopt a new global name - Allergan - pending shareholder approval in 2015.

Forward-Looking State Statements contained in effect Actavis' current praw, Actavis disclaims an Actavis' current expectatine difficulty of predicting market acceptance of an difficulties or delays in market acceptance.	ement this press releaterspective of exity intent or obligations depending the timing or or decontinued demanufacturing; ar	se that refer to future sting trends and in ation to update the upon a number of utcome of FDA approach for Actavis' pand other risks and it	are events or other formation as of the forward-look factors affecting provals or action roducts; risks as uncertainties det	er non-historical factorical factorists of this related as the date of this related as the factorists of the factorists	ease. Except as expetual results may different the competitive products of competitive productions, mergers are riodic public filings.	king statements that ressly required by er materially from ide, among others, ducts and pricing; and joint ventures; with the Securities
and Exchange Commiss 2015. Except as express Actavis Investors: Lisa DeFrancesco (862) 261-7152 Media: David Belian (862) 261-8141	sty required by la	w, Actavis disclain	ns any intent or	obligation to upda	e these forward-loo	king statements.
To view the original vers equining-continued-districtions of the second	ibution-of-name  sk above, please copy of zhunf7s=657788p=kol	and paste the URL below in news&nyo=0	<u>himi</u>		/actavis-contirms-a	

Cli